

# **Higher preoperative serum levels of PD-L1 and B7-H4 are associated with invasive and metastatic potential and predictable for poor response to VEGF-targeted therapy and unfavorable prognosis of renal cell carcinoma**

Takehiko Fukuda <sup>1</sup>, Takao Kamai\* <sup>1</sup>, Akinori Masuda <sup>2</sup>, Akinori Nukui <sup>3</sup>,  
Hideyuki Abe <sup>1</sup>, Kyoko Arai <sup>1</sup>, Ken-Ichiro Yoshida <sup>1</sup>.

<sup>1</sup> Department of Urology, Dokkyo Medical University, Tochigi, Japan.

<sup>2</sup> Dialysis center, Dokkyo Medical University Koshigaya Hospital, Saitama, Japan.

<sup>3</sup> Department of Urology, Nasu Red Cross Hospital, Tochigi, Japan.

T.F. and T.K. are equal contribution to this work

## **\* Address for correspondence**

Takao Kamai M.D.

Department of Urology, Dokkyo Medical University

880 Kitakobayashi Mibu, Tochigi, 321-0293, Japan

Tel: 81-282-87-2162 (international), Fax: 81-282-86-7533

E-mail: [kamait@dokkyomed.ac.jp](mailto:kamait@dokkyomed.ac.jp)

## **A running title**

Availability of serum levels of PD-L1 and B7-H4

## **Novelty and Impact statements**

PD-L1 and B7-H4 are negative regulators of T-cell-mediated antitumor response. Clinical impact of serum levels of the B7 molecular family remains unclear in renal cell

carcinoma (RCC). This study shows that higher preoperative serum levels of PD-L1 and B7-H4 are associated with invasive and metastatic potential of the tumors, a poor response to VEGF-targeted therapy and an unfavorable prognosis, indicating that investigating the preoperative serum levels of B7 family molecules may assist in the design and patient selection strategies of not only VEGF-targeted therapy but also clinical trials of immune checkpoint therapy targeting PD-1/PD-L1 axis.

### **Key Words**

renal cell carcinoma, sunitinib, axitinib, receptor tyrosine kinase inhibitors, immune checkpoints, B7 family, CD28-family receptors

### **Conflict of Interest**

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

### **Grant Support**

This work was supported in part by Japanese Science Progress Society KAKENHI Grants (26462426) to T. Kamai.

### **Abstract**

Renal cell carcinoma (RCC) is an immunogenic and proangiogenic cancer. Although anti-vascular endothelial growth factor (VEGF) therapies achieve impressive responses in some patients, many tumors eventually develop resistance to such therapy. The B7 family molecules such as CTLA-4, PD-1, and PD-L1 are pivotal players in immune checkpoints that positively or negatively regulate various immune responses. Recently, immunotherapy based on blocking immune checkpoints with anti-CTLA4, anti-PD-1, or anti-PD-L1 antibodies has been proposed as a potential new approach to the treatment of

metastatic RCC. Higher expression of PD-L1 and B7-H4 in the tumors is associated with a poor prognosis in RCCs, however, the clinical impact of serum levels of B7 family molecules has not been elucidated in patients with metastatic RCCs receiving VEGF-targeted agents. We assessed the preoperative serum levels of B7 family molecules, including CD80, CD86, PD-1, PD-L1, B7-H3, B7-H4, and CTLA-4, and CD28 in RCC patients, and determined their relations with various clinicopathological characteristics. Elevated preoperative serum levels of PD-L1 and B7-H4 were correlated with less differentiated tumors, higher invasive and metastatic potential, a worse response to anti-VEGF therapy, and shorter overall survival. These findings suggested that investigating preoperative serum levels of PD-L1 and B7-H4 might not only be useful to assess the biological aggressiveness of RCCs, but also to predict the efficacy of anti-VEGF therapy and the eventual prognosis, indicating the future design of clinical trials of therapies targeting immune checkpoint in advanced RCCs.

## **Introduction**

The incidence of renal cell carcinoma (RCC) is steadily increasing and it now accounts for 2-3% of all adult malignancies.<sup>1</sup> About 40% of RCC patients die of metastatic disease, because metastases are often present at diagnosis and relapse following nephrectomy is common.<sup>2</sup> Although localized RCC can be cured by surgery, the prognosis is poor if distant metastases have developed. Approximately 30% of RCC patients have metastatic disease at presentation and recurrence occurs in another 30% of patients after complete resection of the primary tumor.<sup>2</sup>

Clear cell RCC (ccRCC) is a typical immunogenic tumor. It frequently contains high levels of tumor-infiltrating T cells and cytotoxic T cells that recognize and selectively kill tumor cells, and tumor-specific T cells can be detected in the blood of RCC patients.<sup>3-5</sup> Cytokines secreted by tumor cells or host cells play an important role as mediators and/or regulators of various tumor-host interactions involved in the progression of malignancy. T cells have an important role in the host immune response to malignancies, which is most evident in the case of tumors treated by immunotherapy. Cytokines are required for the proliferation and differentiation of effector T cells (Teff) and regulatory T (Treg) cells, and activation and suppression of the immune system show dynamic interaction, with various cytokines exerting effects in either direction and even doing so simultaneously. Thus, while immunocytokine therapy was the mainstay of ccRCC treatment for a long time, its effectiveness was extremely limited.

However, improvements in understanding of the molecular pathways involved have revealed that ccRCC is characterized by high constitutive production of vascular endothelial growth factor (VEGF) that induces a specific tumor vasculature. After the introduction of targeted therapy has ~~dramatically~~ changed the treatment available for metastatic RCC and has improved the progression-free survival of patients by months with this disease.<sup>6</sup> However, many tumors eventually develop resistance to targeted therapy, and only a very small minority of patients remains alive at 5 years due to secondary mutation of the target protein or compensatory changes within the target pathway that bypass the site of inhibition.<sup>7</sup> Accordingly, there has been increasing interest in new immunotherapy modalities that diverge from this paradigm and could potentially improve the outcome of patients with metastatic RCC, including anti-cytotoxic T-lymphocyte-associated antigen 4 (CTLA4) antibody and anti-programmed cell death 1 (PD-1)/PD-ligand 1 (PD-L1) antibody.<sup>8</sup>

Tumor cells need to evade destruction by the immune system to survive and this is now viewed as one of the “hallmarks of cancer”.<sup>9</sup> Recent improvements in our understanding

of tumor progression have shed light on new options for immunotherapy. Important insights into the limitations of T cell-based anticancer immunotherapy have been provided by the discovery of inhibitory pathways termed immune checkpoints (a plethora of inhibitory pathways that are crucial for maintaining self-tolerance and modulating the duration and amplitude of physiological immune responses), and it has been found that tumors co-opt certain immune checkpoint pathways as a major mechanism of immune resistance.<sup>10</sup> Accordingly, the blockade of such immune checkpoints may be a promising approach to the activation of antitumor immunity. The B7 and CD28 families play pivotal roles in the immune checkpoint system by activating and inhibiting co-stimulatory molecules that positively or negatively regulate immune responses.<sup>11,12</sup> Aberrant expression of co-inhibitory molecules from the B7 family has been associated with reduced antitumor immunity and with immune evasion, prompting the development of therapeutic agents that can restore T cell function. Clinical studies investigating the effects of blocking CTLA-4 and PD-1/PD-L1 have yielded exceptionally promising results in patients with advanced melanoma and other cancers, including RCC.<sup>8</sup> However, the role of B7 family molecules in human cancers, including RCC, has been investigated by using tissue samples up to now. Since obtaining blood samples from patients is easier than harvesting tissue by biopsy, circulating biomarkers rather than tissue markers would be useful for assessing the evolution of tumor and immune responses during therapy, as well as for individualized adjustment of treatment. Therefore, we were interested in measuring the serum levels of B7 family molecules and CD28 family receptors in patients with ccRCC. We investigated the preoperative serum levels of B7/CD28 molecules simultaneously in ccRCC patients and to assess correlations with clinicopathological features and the response to treatment. Such information should shed light on the role and biological significance of immune checkpoints in human ccRCCs.

## **Materials and Methods**

## Patients

This is a retrospective study to investigate the preoperative serum levels of B7/CD28 molecules of the 181 patients (109 men and 72 women) in whom ccRCC was diagnosed histopathologically and nephrectomy was performed at our centers between June 2007 and June 2014. Radical nephrectomy for 118 non-metastatic patients (cT<sub>any</sub>N<sub>any</sub>M<sub>0</sub>) or cytoreductive nephrectomy for 63 metastatic patients (cT<sub>any</sub>N<sub>any</sub>M<sub>1</sub>) were performed before any other therapy. All of them underwent preoperative imaging (CT and/or MRI) for tumor staging. The postoperative follow-up period ranged from 3 to 100 months (median: 31 months). The tumor grade and clinical stage were determined according to the Fuhrman system and the TNM classification, respectively.<sup>13,14</sup> In this cohort, the study identified 63 patients with metastatic ccRCC (cT<sub>any</sub>N<sub>any</sub>M<sub>1</sub>) at diagnosed who had undergone cytoreductive nephrectomy and received ~~interferon (IFN)-alpha~~ or tyrosine kinase inhibitors for metastatic lesions as adjuvant therapy. This study was conducted in accordance with the Helsinki Declaration and was approved by the institutional ethical review board of Dokkyo Medical University Hospital. Each patient signed a consent form that was approved by our institutional Committee on Human Rights in Research.

To be included in the study, patients were required to (1) be adults ( $\geq 18$  years old) diagnosed with ccRCC; (2) be treated for 63 metastatic ccRCC at cytoreductive nephrectomy with sunitinib as first-line therapy; (3) have discontinued the first agents for medical reasons (e.g., disease progression, non-response without progression, drug intolerance); (4) have subsequently initiated axitinib as second-line therapy in ; and (5) have their medical records available for review for the entire period encompassing initiation from first therapy until the most recent follow-up or death. Postoperative adjuvant therapy with sunitinib was given to 63 patients (initial dose of 37.5 or 50 mg/day, four weeks on and two weeks off), as first-line systemic therapy after cytoreductive nephrectomy. Within the patients refractory to first-line sunitinib, the 40 patient received axitinib (recommended starting dose of 10 mg/day) as second-line therapy. Progression

was determined by physicians based on radiographic evidence indicating progression of tumor lesions or occurrence of new lesions, physical exams indicating worsening performance status, or cancer-related symptoms (e.g. increased pain, fever and weight loss). Doses of sunitinib and axitinib were decreased if grade 3/4 toxicity occurred. The effect of treatment on the tumors was assessed according to RECIST criteria.<sup>15</sup> The final date of assessment of the disease was collected from patient medical records in September 2015.

Serum samples obtained from the 181 ccRCC patients and 25 healthy controls (14 men and 11 women aged 22-73 years) were stored at  $-80^{\circ}\text{C}$  until analysis. The serum levels of B7 family molecules, CD28, and VEGF were measured by using the following ELISA kits: human CD80 (CSB-E15768h, Cusabio Biotech, China), human CD86 (CSB-E08542h, Cusabio Biotech), human B7-H4 (CSB-E15013h, Cusabio Biotech), human PD-1 (CSB-E13643h, Cusabio Biotech), human PD-L1 (CSB-E13644h, Cusabio Biotech), human CD28 (CSB-E09296h, Cusabio Biotech), human B7-H3 (DB7H30, R&D Systems, MN), human VEGF (DVE00, R&D Systems, MN), and human CTLA-4 (KA0165, Abnova, Taiwan), as described previously.<sup>16</sup> The mean value for three blood samples was used.

### **Statistical analysis**

Baseline characteristics were compared using Wilcoxon rank-sum tests for continuous variables and chi-square tests for categorical variables. Regarding the relationship between preoperative serum levels and variables, statistical analysis was performed by using the Mann-Whitney *U* test to compare two groups and the Kruskal-Wallis test for comparisons among three or more groups. Spearman's rank correlation coefficient analysis was employed to determine the relations between levels of the molecules of interest. Cause-specific survival curves were drawn for the ccRCC patients by the Kaplan-Meier method and differences of survival were examined by the log-rank test. The impact of T cell regulatory molecules, tumor grade, pT stage, pN stage, microscopic

vascular invasion, and distant metastasis (M stage) on survival was assessed by Cox proportional hazards analysis using univariate and multivariate models. In all analyses, a P value < 0.05 was considered to indicate significance. Data were analyzed with commercially available software, as described previously.<sup>16</sup>

## **Results**

### **Preoperative serum levels of B7 family molecules, CD28, and VEGF**

The preoperative serum levels of B7-H4, PD-L1 and VEGF were significantly higher in ccRCC patients than in the healthy controls (Figure 1 and Table 1), but those of PD-1, B7-H3, CTLA-4, CD28, CD80, and CD86 were not (Figure 1, Table 1 and Supplementary Table S1).

Investigation of the correlations between preoperative serum levels of VEGF and B7 family molecules or CD28 family receptors in all cases showed that the preoperative serum VEGF level was closely associated with that of B7-H4 ( $r^2 = 0.66$ , Figure 2A) and weakly related with that of PD-L1 ( $r^2 = 0.39$ , Figure 2C), but not with the levels of PD-1, B7-H3, CTLA-4, CD28, CD80, and CD86. Within metastatic (M1) cases, preoperative serum VEGF level was significantly related with that of B7-H4 ( $r^2 = 0.54$ , Figure 2B) and weakly related with that of PD-L1 ( $r^2 = 0.26$ , Figure 2D).

### **Association of the preoperative serum levels of PD-1, PD-L1, B7-H4, and VEGF with aggressive and metastatic RCC**

The preoperative levels of PD-1, B7-H4, and VEGF were higher in patients with a Karnofsky performance status below 80% than in those above 80% (Table 1).

Higher preoperative serum levels of PD-L1, B7-H4, or VEGF were related to less differentiated tumors, local invasion, regional lymph node metastasis, microscopic vascular invasion, and distant metastasis (Table 1).

An elevated preoperative serum level of PD-1 was correlated with local invasion and microscopic vascular invasion (Table 1).



Elevation of serum CD28 was correlated with local invasion and showed a weaker relation with microscopic vascular invasion (Supplementary Table S1).

Elevation of serum CD80 and CD86 was correlated with less differentiation, local invasion, and microscopic vascular invasion (Supplementary Table S1).

In contrast, serum levels of CTLA-4 and B7-H3 were not correlated with any of these pathological characteristics (Supplementary Table S1).

### **Relationship between serum levels of B7 family molecules and the response to VEGF-targeted therapy for metastatic ccRCC**

Sixty-three patients with metastatic disease at cytoreductive nephrectomy received sunitinib as first-line adjuvant therapy. Of 63 patients, 36 patients had clinical benefit, but 27 patients did not. Eleven patients out of latter 27 patients had best supportive care after sunitinib. Forty patients received axitinib as second-line therapy; 23 patients showed a favorable response, while the other 17 patients did not. The patients with lower preoperative serum levels of PD-L1, B7-H4 and VEGF showed a favorable response to sunitinib or axitinib ( $P=0.0216$ ,  $P=0.0024$ , and  $P=0.0382$ , respectively, Table 1).

In contrast, preoperative serum levels of PD-1, B7-H3, CTLA-4, CD28, CD80, and CD86 were not associated with the response to treatment (Table 1 and Supplementary Table S1).

### **Association of PD-L1 and B7-H4 elevation with shorter overall survival in all ccRCC patients**

The median preoperative serum level of PD-L1 was 18.3, so the patients were divided into two groups at this cut-off value. Kaplan-Meier analysis showed that a higher PD-L1 level was associated with shorter overall survival (Figure 3A). Similarly, a higher B7-H4 level (median: 0.383) and a higher VEGF level (median: 72.6) were also associated with shorter overall survival (Figures 3B,C). The levels of PD-1, B7-H3, CTLA-4, CD28, CD80, and CD86 were not associated with overall survival. While PD-L1, B7-H4, VEGF, less differentiation, local invasion, regional lymph node involvement, microscopic

vascular invasion, and metastasis were significant according to Cox univariate analysis, PD-L1, B7-H4, regional lymph node involvement, and metastasis had impact on shorter overall survival according to multivariate analysis (Table 2).

### **Elevation of PD-L1 and B7-H4 is associated with early relapse after radical nephrectomy in non-metastatic ccRCC patients**

The 108 patients with cN0M0 tumors at nephrectomy were divided into two groups based on the median preoperative serum levels of B7 family molecules, CD28, and VEGF. Comparison of the Kaplan-Meier curves for the low and high value groups revealed that high levels of PD-L1 (median: 14.4), B7-H4 (median: 0.318), and VEGF (median: 41.3) were associated with early relapse after radical nephrectomy (Figure 4), while the other molecules did not show any association. Although less differentiation, local invasion, microscopic vascular invasion, PD-L1, B7-H4, and VEGF were associated with early relapse after nephrectomy according to univariate analysis, only PD-L1 and less differentiation were significant by multivariate analysis (Table 3).

### **Discussion**

One of the most promising approaches to activating therapeutic antitumor immunity is the blockade of immune checkpoints, because it has become clear that tumors utilize these checkpoints as a mechanism of resistance, particularly against T cells specifically targeting tumor antigens.<sup>10</sup> Recent preclinical and clinical studies have demonstrated that checkpoint blockade using anti-CTLA-4, anti-PD-1, and anti-PD-L1 antibodies can be successful for cancer immunotherapy.<sup>18-20</sup> Therefore, it is important to assess whether T cell regulatory molecules can be used as biomarkers to predict tumor resistance and guide the choice of therapy, which is why we investigated the preoperative serum levels of B7 family molecules and CD28 family receptors in RCC patients as well as their associations with clinicopathological features.

There were four main findings of this study. First, the preoperative serum levels of PD-L1, B7-H4, and VEGF in ccRCC patients were significantly higher than those in the healthy controls, and the correlation of B7-H4 or PD-L1 with VEGF in ccRCC patients was established at their higher expression levels, in particular in metastatic disease, indicating that this correlation would have any disease-related implications. Second, the preoperative serum levels of PD-1, PD-L1, B7-H4, and VEGF were correlated with the invasive and metastatic potential of ccRCC. Third, patients with high preoperative serum levels of PD-L1 and B7-H4 showed a poor response to the VEGF-targeted therapy, and had shorter overall survival according to multivariate analysis. Fourth, the higher preoperative serum level of PD-L1 was associated with early relapse after nephrectomy according to multivariate analysis. These findings suggest that the serum levels of molecules in the B7 family might be useful for understanding the biological characteristics of ccRCC and for predicting the response to the VEGF-targeted therapy and the survival.

Increased angiogenesis and immune system evasion are two mechanisms which help ccRCC to proliferate and metastasize. Although VEGF is primarily known for its role in promoting angiogenesis, there is mounting evidence that VEGF also has an immunosuppressive effect and that targeting tumor vasculature may increase the efficacy of immunotherapy. ccRCC can evade immune surveillance by expressing PD-L1, which binds to PD-1 on the surface of activated T and B cells, thus negatively regulating immune activity.<sup>17</sup> There is a strong association between higher tissue levels of PD-L1 expression and adverse clinical outcomes of various cancers, including RCC.<sup>11,21-23</sup> Dong et al. reported that PD-L1 overexpression by tumors *in vivo* and *in vitro* leads to immune tolerance, whereas blockade of PD-L1 or PD-1 enhances antitumor immunity.<sup>24</sup> Blockade of the PD-L1/PD-1 interaction achieves a durable response in approximately 30% of patients with ccRCC, and the response seems to be related to the tumor level of PD-L1 expression.<sup>18,19</sup> However, it remains challenging to select patients who are likely

to respond to anti-PD-1/PD-L1 therapy.<sup>12</sup> In the COMPARZ trial, RCC patients received sunitinib or pazopanib therapy and increased immunohistochemical tumor cell PD-L1 expression was associated with shorter survival.<sup>25</sup> In the present study, higher preoperative serum level of PD-L1 was correlated with poorer differentiation, invasive and metastasis profiles, and shorter overall survival, and was independent prognostic factors for overall survival. Frigola et al. reported that an elevated preoperative level of soluble PD-L1 is associated with an increased risk of death from ccRCC, and they suggested that release of soluble PD-L1 molecules may be a mechanism by which tumors impair systemic antitumor immunity, suggesting that inactivation or removal of soluble PD-L1 from the serum could be clinically beneficial.<sup>26</sup> Taken together, these findings have focused attention on blockade of PD-1/PD-L1 as a possible strategy for antitumor immunotherapy in patients with advanced RCC.

B7-H4 is a negative regulatory molecule expressed on the cell membrane that inhibits proliferation and cytokine production by CD4 and CD8 T cells.<sup>11</sup> In general, B7-H4 positivity is associated with advanced cancer and poor survival, and this is also the case for RCC.<sup>11,27</sup> In an experimental model of lung metastasis, B7-H4 expression by host cells was demonstrated to suppress tumor-specific T cell responses and promote the infiltration of immunosuppressive T cell subsets into the lungs. As a result, B7-H4 WT mice had more lung metastases than B7-H4 knockout mice.<sup>28</sup> Expression of B7-H4 was also reported to show a correlation with tumor infiltration by Treg cells.<sup>29</sup> Tumor-infiltrating lymphocytes may be a manifestation of antitumor immunity, but increased tumor infiltration by T cells is associated with shorter survival of RCC patients,<sup>4</sup> indicating that there is a potential for the mechanisms of T cell immunity to fail in RCC. Zhang et al. recently reported that B7-H4 contains a functional nuclear localization sequence (NLS), which is responsible for the nuclear localization of this molecule in human RCC tissues.<sup>30</sup> They also reported that a point mutation of the B7-H4 NLS motif suppressed leptomyacin B - induced nuclear accumulation of B7-H4 and abrogated

B7-H4-mediated proliferation and cell cycle progression of HEK293 human embryonic kidney cells, while wild-type B7-H4 confers chemoresistance on RCC cell lines, indicating that nuclear localization of B7-H4 might be crucial for its influence on cell proliferation and cell cycle progression.<sup>30</sup> Thompson et al. reported in 101 human RCC patients that the higher serum-soluble B7-H4 levels were associated with poorer differentiation, local invasion, lymph node metastasis and distant metastasis.<sup>31</sup> Zhang et al. reported in 93 human hepatocellular carcinoma (HCC) patients that soluble B7-H4 levels in patients were significantly higher than that in healthy controls; high sB7-H4 levels were correlated with tumor differentiation, tumor invasion, tumor-node metastasis stage, and poor overall survival; high sB7-H4 levels were independent prognostic factors for overall survival.<sup>32</sup> These findings are consistent with our observation in RCCs in this study. Thus, serum levels of B7-H4 may be potential diagnostic and prognostic marker. Taken together, these observations provide a basis for targeting B7-H4 in cancer and B7-H4 is thus believed to have potential as a biomarker. Furthermore, given the Cox multivariate analyses in which PD-L1, B7-H4, regional lymph node involvement, and metastasis had significant impact on shorter overall survival, it might be possible to provide a better prognostic model using PD-L1, B7-H4 with either lymph node status or metastatic status for ccRCC patients. Similarly, since PD-L1 and differentiation were significant for recurrence-free survival, preoperative serum level of PD-L1 and histological grade might be available for predicting recurrence in N0M0 cases after nephrectomy.

VEGF is a master regulator of angiogenesis, and also has profound effects on immune regulatory cell function, specifically inhibiting dendritic cells (DC) maturation and antigen presentation. For many tumors, hypoxia causes tumor cells, stromal cells, and normal cells to produce VEGF with immunosuppressive effects both locally and at distant sites, and creates a microenvironment of inhibitory inflammatory cells; the

recruitment of myeloid-derived suppressor cells (MDSC) and the induction of Treg cells by inhibiting DC maturation.<sup>33-35</sup>

On the other hand, it is not clear that PD-L1, B7-H4 and VEGF could evade immune surveillance in the cooperative or independent manner by negatively regulating the immune system in ccRCC, and whether VEGF is the causal factor in release of PD-L1 or B7-H4 expressed by RCC to peripheral blood. In the present study, the preoperative serum VEGF level was associated with that of PD-L1 and B7-H4, and higher preoperative serum levels of PD-L1, B7-H4, or VEGF were related to less differentiated tumors, local invasion, regional lymph node metastasis, microscopic vascular invasion, and distant metastasis. Therefore, we hypothesize that tumors with increased VEGF expression would possess an immunosuppressive tumor microenvironment and tumor-derived PD-L1, B7-H4, and VEGF may locally inactivate immune cells. However, as Frigola et al. suggested, systemic effects either by recirculation of immune cells through PD-L1, B7-H4, and VEGF-positive tumor sites or by the release of biologically active soluble forms of PD-L1, B7-H4, and VEGF into the circulation cannot be excluded, indicating that both scenarios can contribute to global immunosuppression.<sup>26</sup>

In contrast, the preoperative serum levels of CTLA-4, B7-H3, CD28, CD80, and CD86 were not correlated with tumor biological characteristics in the present study. CTLA-4 is a member of the immunoglobulin gene superfamily and is a B7-binding protein along with its homologue (CD28).<sup>11</sup> It plays an important role in downregulation of T cell responses, T cell homeostasis, and maintenance of peripheral tolerance. A CTLA-4 antibody was the first member of this class of immunotherapy agents to receive US FDA approval, and it achieved promising results in a recent phase II study of metastatic ccRCC.<sup>11</sup> CD28 binds CD80 and CD86 expressed by activated antigen-presenting cells and plays as the prototypic T cell costimulatory receptor that augments TCR-mediated T cell proliferation, survival, metabolic activity, and effector function.<sup>10,11</sup> CD80 and CD86 are homologous costimulatory molecules predominantly expressed on antigen-presenting

cells. Interaction of these B7 family molecules with CD28 and CTLA-4 expressed on T cells is a critical step in T cell activation. Despite their structural similarities, CD80 and CD86 make distinct contributions to the development of pathogenic T cells in autoimmune diseases and protective T cells in infectious diseases. While B3-H7 shows structural similarities to the classical co-stimulatory molecules CD80 and CD86, there have been contradictory findings regarding its possible co-stimulatory or co-inhibitory role. Thus, the relationship between B7-H3 expression and the prognosis of cancer patients remains to be elucidated.

The limitations of our study are retrospective design and the present study investigated a relatively small number of patients and the follow-up period was too short to draw definite conclusions. The findings of our study need further validation in forthcoming studies, better in prospective controlled large sampled clinical trials. As biomarkers may not be static, serial measurement may be required to assess the evolution of tumor and immune responses during therapy, as well as to adjust treatment in a personalized manner. In this context, in addition to examination of the expression of PD-L1, B7-H4 and VEGF in the primary tumors, repeated measurement of the serum levels of B7 family molecules may be important for monitoring both tumor behavior and the condition of patients in the prospective studies on a larger scale. If we accept the hypothesis that release of soluble T cell suppressor molecules is a mechanism by which tumors interfere with systemic antitumor immunity, it is also important to determine whether the serum levels of soluble T cell regulatory molecules (such as PD-L1, B7-H4, PD-1, and CTLA-4) can be used as biomarkers to predict the effectiveness of immunotherapy targeting T cells in patients with ccRCC. Further investigation is needed to evaluate the mechanistic basis of the ability to counteract the dual function of angiogenic factors in promoting vessel growth and in suppressing immune responses by the parallel effects on host immunity and tumor vasculature in ccRCC. It is hoped that further investigation of the role of T cell regulatory

molecules in promoting tumor immune tolerance will lead to the development of new immunotherapy options for ccRCC.

### **Acknowledgements**

The authors thank all patients and their families for contribution to this study.

### **References**

1. Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin.* 2012;62:10–29.
2. Rini B, Campbell SC, Escudier B. Renal cell carcinoma. *Lancet.* 2009;373:1119-1132.
3. Bromwich EJ, McArdle PA, Canna K, et al. The relationship between T-lymphocyte infiltration, stage, tumour grade and survival in patients undergoing curative surgery for renal cell cancer. *Br J Cancer.* 2003;89:1906–1908.



4. Nakano O, Sato M, Naito Y, et al. Proliferative activity of intratumoral CD8(+) T-lymphocytes as a prognostic factor in human renal cell carcinoma: clinicopathologic demonstration of antitumor immunity. *Cancer Res.* 2001;61:5132–5136.
5. Jantzer P, Schendel DJ. Human renal carcinoma antigen-specific CTLs, antigen-driven selection and long-term persistence *in vivo*. *Cancer Res.* 1998;58:3078-3086.
6. Abe H, Kamai T. Recent advances in the treatment of metastatic renal cell carcinoma. *Int J Urol.* 2013;20:944-955.
7. Vanneman M, Dranoff G. Combining immunotherapy and targeted therapies in cancer treatment. *Nat Rev Cancer.* 2012;12:237-251.
8. Harshman LC, Drake CG, Wargo JA, Sharma P, Bhardwaj N. Cancer immunotherapy highlights from the 2014 ASCO meeting. *Cancer Immunol Res.* 2014; 2:714-719.
9. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell.* 2011;144:646-674.
10. Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer.* 2012;12:252-264.
11. Zou W, Chen I. Inhibitory B7-family molecules in the tumour microenvironment. *Nat Rev Immune.* 2008;8:467-477.
12. Ribas A, Tumei PC. The Future of Cancer Therapy: Selecting Patients Likely to Respond to PD1/L1 Blockade. *Clin Cancer Res.* 2014;20:4982-4498.
13. Fuhrman SA, Lasky LC, Lmas C. Prognostic significance of morphologic parameters in renal cell carcinoma. *Am J Surg Pathol.* 1982;6:655-663.
14. Sobin LH, Gospodarowicz MK, Wittekind CH editors. International union against cancer. UICC, In TNM classification of malignant tumors, 7th ed. New York, Wiley-Liss, 2009, 255-257.
15. Therasse P, Arbuck SG, Eisenhauer EA, et al. New guidelines to evaluate the response to treatment in solid tumors. European Organization for research and Treatment

of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst.* 2000; 92:205–216.

16. Masuda A, Arai K, Nishihara D, et al. Clinical significance of serum soluble T cell regulatory molecules in clear cell renal cell carcinoma. *Biomed Res Int.* 2014; 2014: 396064.

17. Kim H-J, Cantor H. The Path to Reactivation of Antitumor Immunity and Checkpoint Immunotherapy. *Cancer Immunol Res.* 2014;2:926-936.

18. Topalian SL, Hodi FS, Brahmer JR, et al. Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med.* 2012;366:2443–2454.

19. Brahmer JR, Tykodi SS, Chow LQ, et al. Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *N Engl J Med.* 2012;366:2455–2465.

20. Gunturi A, McDermott DF. Potential of new therapies like anti-PD1 in kidney cancer. *Curr Treat Options Oncol.* 2014;15:137-146.

21. Thompson RH, Gillett MD, Cheville JC, et al. Costimulatory B7-H1 in renal cell carcinoma patients: indicator of tumor aggressiveness and potential therapeutic target. *Proc Natl Acad Sci USA.* 2004;101:17174–17179.

22. Krambeck AE, Dong H, Thompson RH, et al. Survivin and b7-h1 are collaborative predictors of survival and represent potential therapeutic targets for patients with renal cell carcinoma. *Clin Cancer Res.* 2007;13: 1749–1756.

23. Thompson RH, Kuntz SM, Leibovich BC, et al. Tumor B7-H1 is associated with poor prognosis in renal cell carcinoma patients with long-term follow-up. *Cancer Res.* 2006; 66: 3381–3385.

24. Dong H, Strome SE, Salomao DR, et al. Tumor-associated B7-H1 promotes T-cell apoptosis: a potential mechanism of immune evasion. *Nat Med.* 2002;8:793-800.

25. Choueiri TK, Figueroa DJ, Fay AP, et al. Correlation of PD-L1 tumor expression and treatment outcomes in patients with renal cell carcinoma receiving sunitinib or

- pazopanib: Results from COMPARZ, a randomized controlled trial. *Clin Cancer Res.* 2014; 21: 1071-1077.
26. Frigola X, Inman BA, Lohse CM, et al. Identification of a soluble form of B7-H1 that retains immunosuppressive activity and is associated with aggressive renal cell carcinoma. *Clin Cancer Res.* 2011; 17: 1915-1923.
27. Krambeck AE, Thompson RH, Dong H, et al. B7-H4 expression in renal cell carcinoma and tumor vasculature: associations with cancer progression and survival. *Proc Natl Acad Sci USA.* 2006; 103: 10391-10396.
28. Abadi YM, Jeon H, Ohaegbulam KC, et al. Host b7x promotes pulmonary metastasis of breast cancer. *J Immunol.* 2013; 190: 3806–3814.
29. Kryczek I, Wei S, Zhu G, et al. Relationship between B7-H4, regulatory T cells, and patient outcome in human ovarian carcinoma. *Cancer Res.* 2007; 67: 8900–8905.
30. Zhang L, Wu H, Lu D, et al. The costimulatory molecule B7-H4 promote tumor progression and cell proliferation through translocating into nucleus. *Oncogene.* 2013; 32: 5347-5358.
31. Thompson RH1, Zang X, Lohse CM, et al. Serum-soluble B7x is elevated in renal cell carcinoma patients and is associated with advanced stage. *Cancer Res.* 2008; 68: 6054-6058.
32. Zhang C, Li Y, Wang Y. Diagnostic value of serum B7-H4 for hepatocellular carcinoma. *J Surg Res.* 2015; 197: 301-306.
33. Huang Y, Goel S, Duda DG, Fukumura D, Jain RK. Vascular normalization as an emerging strategy to enhance cancer immunotherapy. *Cancer Res.* 2013;73:2943–8.
34. Gabrilovich DI, Ostrand-Rosenberg S, Bronte V. Coordinated regulation of myeloid cells by tumours. *Nat Rev Immunol.* 2012; 12: 253–268.
35. Tartour E, Pere H, Maillere B, Terme M, Merillon N, Taieb J, et al. Angiogenesis and immunity: a bidirectional link potentially relevant for the monitoring of antiangiogenic

therapy and the development of novel therapeutic combination with immunotherapy. *Cancer Metastasis Rev* 2011;30:83–95.

## **Figure Legends**

### **Figure 1: Preoperative serum levels between healthy volunteers and RCC patients.**

Preoperative serum levels of VEGF (A), B7-H4 (B) and PD-L1 (C) were higher in RCC patients than those in healthy volunteers, while those of PD-1 (D) were not. The median value is the central line, the box is the interquartile range, the bars are the full range, and the points are the outliers. Bold circled P values were obtained by comparing the two groups with the Mann-Whitney *U* test.

### **Figure 2: Relationship between serum B7 family molecules.**

Spearman rank correlation between the preoperative serum levels of VEGF and B7-H4/PD-L1 in all 171 cases (A, C) and in 83 metastatic (M1) cases at nephrectomy (B, D). The preoperative serum levels of B7-H4 and PD-1 were positively associated with those of VEGF in all case and in M1 cases.

### **Figure 3: Overall survival in all 181 cases.**

This survival curve is based on the median values of preoperative serum levels of B7 family molecules in all 181 cases. The cases were divided into two groups at this level - high and low expression. *P* value was analyzed by log-rank test. The patients with higher preoperative serum levels of PD-L1 (A), B7-H4 (B) and VEGF (C) had poorer overall survival.

**Figure 4: Recurrence free-survival in 108 N0M0 cases at nephrectomy.**

This survival curve is based on the median values of preoperative serum levels of B7 family molecules in non-metastatic (M0) cases at nephrectomy. The cases were divided into two groups at this level - high and low expression. *P* value was analyzed by log-rank test. Higher preoperative serum levels of PD-L1 (A), B7-H4 (B) and VEGF (C) in N0M0 patients at nephrectomy were associated with shorter recurrence free- survival.