

Original

Relationships between Sensory Processing, Aberrant Behaviors and Food-related Behaviors in Individuals with Prader-Willi Syndrome

Asami Takahashi, MD^{1,2*}, Hiroshi Ihara, MD, PhD^{1*}, Hiroyuki Ogata, MA^{1,3},
Masao Gito, MD^{1,2,3}, Nobuyuki Murakami, MD, PhD⁴, Yuji Oto, MD, PhD⁴,
Atsushi Ishii, MD^{1,5}, Sohei Saima¹, MD, Toshiro Nagai, MD, PhD⁶

*Equally contributed to this work.

¹ Department of Psychiatry, Dokkyo Medical University Saitama Medical Center, Saitama, Japan

² Ikezawa Hospital, Saitama, Japan

³ Department of Psychiatry, Dokkyo Medical University School of Medicine, Tochigi, Japan

⁴ Department of Pediatrics, Dokkyo Medical University Saitama Medical Center, Saitama, Japan

⁵ Department of Psychiatry, National Hospital Organization Hanamaki Hospital, Iwate, Japan

⁶ Nakagawanosato Ryoiku Center, Saitama, Japan

SUMMARY

Objective : The level of sensory processing dysfunction was examined and compared with the severity of food-related behaviors and aberrant behaviors in 102 individuals (60 males and 42 females) with Prader-Willi syndrome (PWS), including 76 patients with paternally inherited deletion and 26 patients with maternal uniparental disomy within the 15q11-13 region.

Methods : The Japanese version of the Short Sensory Profile (SSP-J), the Food Related Problem Questionnaire (FRPQ) and Aberrant Behavior Checklist Japanese Version (ABC-J) were administered to PWS patients.

Results : Based on the results of SSP-J, the patients in this study were classified as follows : 27 individuals with Typical Performance, 45 with Probable Difference and 30 with Definite Difference. Among the three groups, one-way analyses of variance (ANOVAs) were conducted to investigate differences in scores of FRPQ and ABC-J. No significant differences were found in the total scores and three subscores of FRPQ. On the contrary, statistically significant differences were found in the total score as well as five subscores in the ABC-J. Post-hoc Tukey's tests revealed significant differences of aberrant behaviors in the total score and all of the five subscores of the ABC-J. Definite Difference was more than Probable Difference and Typical Performance.

Conclusions : Approximately three quarters of individuals with PWS demonstrated abnormalities in sensory responsiveness. In terms of the relationships of sensory processing with other behavioral symptoms,

aberrant behaviors and food-related problems brought out a sharp contrast : the former showed significant associations with sensory processing, whereas the latter did not.

Key Words : Prader-Willi syndrome, sensory profile, sensory processing

Received January 28, 2019 ; accepted February 14, 2019

Reprint requests to : Hiroshi Ihara, MD, PhD

Department of Psychiatry, Dokkyo Medical University Saitama Medical Center, 2-1-50 Minami-Koshigaya, Koshigaya City, Saitama 343-8555 Japan.

INTRODUCTION

Sensory processing means the modulation response of an individual to detect and integrate sensations from multiple sensory systems (e.g., auditory, visual, somatosensory, tactile, taste, smell). The ability of this function is essential in allowing the individual to regulate the flood of sensory stimulation from different modalities^{1,2)}. To modulate sensory stimuli in an appropriate manner is vital for an individual to exclude noise stimuli and to pick out targets from the myriads of sensory inputs. Without proper functioning of sensory modulation, the individual is overwhelmed by too much stimulation and is unable to understand meaningful information from his or her perceptual experiences.

The relationship of sensory processing impairments with behavioral symptoms has been studied in a variety of developmental disorders : childhood autism³⁾, attention-deficit hyperactivity disorder (ADHD)⁴⁾, fragile X syndrome⁵⁾, and Williams syndrome^{2,6)}. So far, four types of sensory processing tendencies have been postulated. The first type is sensory over-responsiveness in which the individual responds to sensory stimuli more vividly than usual. The second one is under-responsiveness that occurs when the individual does not react as quickly compared to usual and/or even disregards sensory stimuli. The third one is sensory seeking behavior in which the individual purposely seeks intense sensory experiences. The last one is sensation avoiding⁷⁾. It has been argued that problematic behaviors associated with various developmental conditions are potentially derived secondarily from the effect of underpinning impairments in sensory modulation. Therefore, the relationship between sensory processing dysfunctions and other behavioral symptoms is worth investigating in a variety of neurodevelopmental disorders.

Prader-Willi syndrome (PWS) is one of such neurodevelopmental disorders, in which there is paucity of research with respect to sensory processing disorders associated with behavioral symptoms. PWS is characterized by the four main features : neonatal hypotonia, intellectual disability, hyperphagia, progressive obesity and hypogonadism^{8,9)}. The physical manifestations of PWS include : short stature, small hands and feet,

hypopigmentation and craniofacial anomalies. As a contiguous gene syndrome, PWS is attributed to a loss of expression of the paternally derived genes in the q11-13 region of chromosome 15. There are mainly two origins of the loss : a paternal deletion (DEL) of 15q11-13 found in 70% of patients, and maternal uniparental disomy 15 (mUPD ; when both copies of chromosome 15 are maternally inherited) found in 25%¹⁰⁻¹³⁾. The remaining 1-3% of mechanism is an imprinting defect (ID), which means a defect in the genomic region that controls the activity of imprinted genes.

Behavioral characteristics of this syndrome have been well studied. Those include food-related behaviors¹⁴⁾, temper outbursts¹⁵⁾, compulsive and ritualistic behaviors¹⁶⁻¹⁸⁾, excoriating behaviors^{19,20)} and autistic-like behaviors^{21,22)}. Ample evidence shows that the mUPD subtype has a higher risk for autistic-like behaviors than the DEL subtype²³⁻²⁵⁾. Such findings in terms of the affinity of mUPD with autism spectrum disorder (ASD) suggest the existence of maternally active gene (s) in chromosome 15q11-13^{23,26-28)}.

Sensory processing abilities in PWS have not been examined yet, while those in ASD have been well researched. A number of studies have shown that ASD have atypical responses to a variety of sensory modalities²⁹⁻³⁴⁾. These findings have been observed in individuals with high functioning ASD³⁵⁾. For diagnostic purpose, sensory processing impairments were not included in the three core behavioral deficits : social relatedness, communication skills and the presence of stereotyped behavior. Eventually, they were listed as parts of criterion B termed 'RESTRICTED, REPETITIVE PATTERNS OF BEHAVIOR, INTERESTS, OR ACTIVITIES AS MANIFESTED BY AT LEAST' in the newest edition of the DSM-5³⁶⁾. Nevertheless, sensory modulation difficulties can be the first noticeable signs for parents to recognize in their children with ASD. As Ben-Sasson et al.³⁷⁾ confirmed, the early emergence of sensory processing dysfunction in toddlers often indicates that such disorder grows to influence a child's adaptive behaviors from an early stage of development. Individuals with PWS that have sensory processing difficulties would imply the possibility of a positive relationship between sensory modulation impairment and the severity of social maladaptive

Table 1 Participants Characteristics

	total	SSP-J total			<i>p</i> -value
		Typical	Probable Difference	Definite Difference	
Number	102	27	45	30	
DEL/mUPD	76/26	19/8	35/10	22/8	
Male/Female	60/42	13/14	30/15	17/13	
<18y.o./18y.o. ≤	52/50	16/11	23/22	13/17	
Mean age	17.98	16.63	17.96	19.23	0.55
Age range	6-48	8-28	6-46	7-48	
IQ mean (N)	48.69 (73)	48.63 (19)	48.77 (31)	48.64 (23)	1.00
IQ range	39-84	39-84	39-79	39-76	

p-values from the one-way analysis of variance

behaviors in this syndrome.

For the first time, this study will survey the broad range of sensory processing ability in individuals with PWS. This study has mainly three objectives. The first is to explore the overall picture of sensory profiles in PWS and to illuminate how frequently the sensory profile differences occur in individuals with PWS and which domains of sensory processing (e.g. tactile, taste/smell, movement, auditory) are considered impaired. The second is to examine the differences between DEL and mUPD and those between female and male in terms of sensory processing impairments. Lastly, the level of sensory processing dysfunction is to be compared with the severity of other behavioral symptoms, such as food-related behaviors and aberrant behaviors.

SUBJECTS AND METHODS

Before starting this study, the Institutional Review Board of Dokkyo Medical University assessed and approved that all procedures conformed the World Medical Association Declaration of Helsinki (No. 21107). Informed consents for behavioral and psychiatric assessment and those specific for cytogenetic and/or molecular-genetic studies were obtained from participants or their parents.

Subjects

This study enrolled 102 Japanese participants with PWS recruited from the Departments of Pediatrics and Psychiatry, Dokkyo Medical University Saitama Medical Center. All patients were diagnosed with

PWS using fluorescence in situ hybridization or the methylation test. The participants consisted of 60 male individuals and 42 female individuals, including 76 patients confirmed to having a DEL involving 15q11-13, and 26 patients confirmed to having mUPD of chromosome 15 (Table 1).

Methods

The Assessment of Behavior

Before the assessment of behavior, a Japanese version of the Wechsler Intelligence Scale³⁸⁻⁴¹⁾ was administered for the measurement of IQ (WISC-III, WAIS-III).

A comprehensive set of behavioral assessment was used in terms of sensory profile, food-related problems, and aberrant behaviors. The psychologist (H.O.) who took data was not informed of the genetic status of each patient. In order to complete data taking, HO had 3 to 6 sessions for each participant. Behavioral assessments applied in this study were originally constructed on the assumption that they were applied as self-administered or informant-based scale. However, some parts of the questionnaire instructions are difficult to understand for participants and parents. Hence, H.O. administered all behavioral measures in face-to-face interview of the individuals or the parents of the PWS patients, immediately before checking for completeness and accuracy. In consequence, the quality of data gathered in this study was expected to be better than that gathered by means of mail-out survey of a questionnaire battery.

Sensory Profile

Sensory processing ability of all participants were examined by means of the Japanese version of the Short Sensory Profile (SSP-J)⁴²⁾. SSP-J consisted of 38 questions, caregivers were asked to grade the frequency that their child showed sensory processing behaviors on the basis of a five-point Likert scale (*always, frequently, occasionally, seldom, or never*)⁴³⁾. The questionnaire included seven subscores: Tactile Sensitivity, Taste/Smell Sensitivity, Movement Sensitivity, Underresponsive/Seeks Sensation, Auditory Filtering, Low Energy/Weak, and Visual/Auditory Sensitivity. A higher total score meant a more severe impairment. The SSP-J has been widely used in Japan. Its internal consistency reliability of each section, including the 7 subscores and total score, in 1441 typically developing children in Japan was between 0.69 and 0.84.

The raw scores of 8 sections were converted to standardized z-scores based on the criteria proposed by McIntosh et al.⁴⁴⁾. In the child's responses to sensory experiences, Typical Performance was defined as z-scores above -1.00, Probable Difference as those from -1.00 to -2.00, and Definite Difference as those below -2.00.

Food-related behaviors

The severity of food-related behaviors was assessed by the means of the Food Related Problem Questionnaire (FRPQ). This informant-based questionnaire was constructed exclusively for the purpose of evaluating the level of eating behaviors in individuals with PWS. FRPQ consisted of 16 items, which are divided into three subscales: preoccupation with food (P), impairment of satiety (S), and other food-related negative behaviors (N). Examples of the questions include: "How often does the person compare the size or content of their meal with others?" (P); "After a normal sized meal, how often does the person say they still feel hungry?" (S); and "If given the opportunity, how often would the person 'help themselves' to food which they should not have?" (N). Data was presented to show that the FRPQ has reliable psychometric properties to appraise the food-related problems in individuals with PWS⁴⁵⁾.

Aberrant Behaviors

In order to examine the level of autistic-like repetitive behaviors and other maladaptive behaviors, the Aberrant Behavior Checklist Japanese Version (ABC-J)⁴⁶⁾ was administered to all participants. This consisted of a 58-item checklist that took about 10-15 minutes to fill in. All items were classified into five categories: a) irritability and agitation, b) lethargy and social withdrawal, c) stereotypic behavior, d) hyperactivity and noncompliance, and e) inappropriate speech. The ABC was found to be an effective tool to identify behavioral manifestations in individuals with intellectual disability⁴⁷⁾ and autism spectrum disorder⁴⁸⁾. This tool was also used for the purpose of measuring treatment response^{47,49)}.

RESULTS

Descriptive statistics: sensory processing difference

Based on the results of SSP-J, the participants in this study were classified as follows: 27 individuals with Typical Performance, 45 with Probable Difference and 30 with Definite Difference (Table 1). One-way analyses of variance (ANOVAs) revealed no significant differences in terms of age and IQ among these three groups.

As Table 2 shows, thirty (29.4%) individuals with PWS demonstrated clinically definite dysfunction of sensory processing. The most prominent features were found in Low Energy/Weak section (49.0% in Definite Difference and 41.2% in Probable Difference). Equally pronounced was the fact that more than half of the individuals were rated as Definite or Probable Difference in Tactile Sensitivity, Movement Sensitivity and Underresponsive/Seeks Sensation. On the other hand, profound impairment was not observed in Taste/Smell Sensitivity, Auditory Filtering and Visual/Auditory Sensitivity, for more than 60% individuals were classified as Typical Performance in these categories.

Genotypical and gender differences

T-tests were conducted to examine the differences between DEL and mUPD in terms of the raw scores of 8 sections. As Table 3 shows, significant differences were not found in any of these sections. A marginal difference was found in 'Auditory Filtering', in which

Table 2 The number of PWS individuals in each three SSP-J total score groups

	SSP-J		
	Typical	Probable Difference	Definite Difference
SSP-J total score	27 (26.5%)	45 (44.1%)	30 (29.4%)
Subscore			
Tactile Snsitivity	40 (39.2%)	52 (51%)	10 (9.8%)
Taste/Smell Sensitivity	83 (81.4%)	19 (18.6%)	0 (0%)
Movement Sensitivity	43 (42.2%)	40 (39.2%)	19 (18.6%)
Underresponsive/Seeks Sensation	47 (46.1%)	42 (41.2%)	13 (12.7%)
Auditory Filtering	66 (64.7%)	30 (29.4%)	6 (5.9%)
Low Energy/Weak	10 (9.8%)	42 (41.2%)	50 (49.0%)
Visual/Auditory Sensitivity	78 (76.5%)	16 (15.7%)	8 (7.8%)

Table 3 SSP-J scores and the results of t-test between DEL and mUPD, and that between male and female.

	genotype		gender		<i>p</i> -value	
	DEL mean ± SD	mUPD mean ± SD	male mean ± SD	female mean ± SD	genotype	gender
SSP-J total score	68.52 ± 18.09	71.23 ± 19.99	69.87 ± 18.37	67.67 ± 19.14	0.52	0.56
Subscore						
Tactile Snsitivity	11.37 ± 3.76	12.19 ± 5.98	11.82 ± 4.41	11.14 ± 4.46	0.52	0.45
Taste/Smell Sensitivity	4.33 ± 0.78	4.5 ± 0.91	4.32 ± 0.7	4.45 ± 0.94	0.37	0.41
Movement Sensitivity	5.77 ± 2.69	5.62 ± 2.52	5.52 ± 2.57	5.98 ± 2.73	0.79	0.39
Underresponsive/Seeks Sensation	11.04 ± 4.22	11.96 ± 4.1	11.98 ± 4.57	10.17 ± 3.35	0.34	0.02*
Auditory Filtering	10.41 ± 4.1	12.23 ± 4.41	10.98 ± 4.33	10.62 ± 4.41	0.06	0.67
Low Energy/Weak	18.75 ± 6.73	17.77 ± 6	18.28 ± 6.52	18.62 ± 6.66	0.51	0.80
Visual/Auditory Sensitivity	6.97 ± 2.7	6.96 ± 2.44	7.13 ± 2.75	6.69 ± 2.42	0.98	0.40

**p*<0.05

mUPD showed a slight trend of impairment (*p*=0.06). Examining gender differences with respect to sensory processing impairments, t-tests were applied in terms of the raw scores of 8 sections. Understandably there were no differences in the total score and six among the seven subscores of SSP-J. In regards to the section of 'Underresponsive/Seek Sensation', an exceptional gender difference was observed in that males showed more severe impairment than their female counterparts (*p*=0.02).

Sensory processing and food-related behaviors

To compare the level of sensory processing with the severity of food-related problems in PWS, ANOVAs were conducted to investigate differences in scores of FRPQ among three groups classified based on the SSP-J results : Typical Performance, Probable Difference and Definite Difference. No significant dif-

ferences were found in the total scores and three subscores of FRPQ, such as preoccupation with food (P), impairment of satiety (S) and other food-related negative behaviors (N) (Table 4).

Sensory processing and aberrant behaviors

For assessing the relationship between sensory processing and aberrant behaviors, one-way ANOVAs were used to examine scores of ABC-J among the three groups. Statistically significant differences were found in the total score as well as five subscores in the ABC-J (Table 4). The five subscores included excitement, apathy, stereotype, hyperactivity and inappropriate speech. In all scores of ABC-J, the individuals with Definite Difference in terms of sensory processing showed the most severe impairment in aberrant behaviors. Those with Probable Difference came in second, and those with Typical Performance

Table 4 FRPQ and ABC-J scores and the results of one-way ANOVA among the three SSP-J total score groups.

	Total	SSP-J total score			ANOVA interaction	
		Typical (N = 27)	Probable Difference (N = 45)	Definite Difference (N = 30)	F	<i>p</i>
FRPQ total score	37.46 ± 14.54	36.00 ± 14.96	38.91 ± 12.30	36.70 ± 17.21	0.39	0.68
Subscore						
FRPQ-P	10.17 ± 4.10	9.44 ± 4.53	10.67 ± 3.46	10.10 ± 4.57	0.75	0.48
FRPQ-S	16.41 ± 5.89	16.52 ± 6.06	16.77 ± 4.56	15.80 ± 7.37	0.24	0.79
FRPQ-N	10.79 ± 6.64	10.04 ± 6.72	11.26 ± 6.14	10.80 ± 7.39	0.28	0.76
ABC-J total score	34.41 ± 30.10	19.19 ± 22.42	29.20 ± 27.30	55.73 ± 28.98	14.88	<0.01**
Subscore						
ABC-J irritability and agitation	11.59 ± 10.07	6.81 ± 6.88	9.93 ± 9.54	18.33 ± 10.00	12.82	<0.01**
ABC-J lethargy and social withdrawal	7.87 ± 10.01	4.11 ± 8.92	6.73 ± 10.89	12.93 ± 7.50	6.72	<0.01**
ABC-J stereotypic behavior	2.24 ± 3.26	0.89 ± 1.95	1.82 ± 2.83	4.07 ± 3.99	8.51	<0.01**
ABC-J hyperactivity and noncompliance	8.82 ± 9.34	4.56 ± 6.15	7.86 ± 9.47	14.07 ± 9.33	9.03	<0.01**
ABC-J inappropriate speech	4.58 ± 3.33	2.78 ± 2.47	4.50 ± 3.39	6.33 ± 3.09	9.49	<0.01**

* $p < 0.05$, ** $p < 0.01$.

FRPQ-P : preoccupation with food, FRPQ-S : impairment of satiety, FRPQ-N : composite negative behavior.

showed the least. Post-hoc Tukey's test revealed significant differences of aberrant behaviors as follows. In ABC-J total score and all of the five scores, Definite Difference was more than Probable Difference and Typical Performance (Fig. 1).

DISCUSSION

This study illuminated that approximately three quarters of individuals with PWS demonstrated abnormalities in sensory responsiveness, as only 26.5% of the entire sample were classed as Typical Performance on the basis of the total score of the SSP-J. When seven subscores of the SSP-J were probed, the most striking feature in PWS individuals was the severe abnormality in Low Energy/Weak section, in which only 9.8% of the sample in this study were classified into Typical Performance. Even in the cases of ASD, 58.0% of the individuals showed Typical Performance in the data of Tomchek and Dunn⁵⁰⁾, who applied the Short Sensory Profile (SSP) to 281 children with ASD. In PWS, equally severe impairment was found in Tactile Sensitivity, Movement Sensitivity and Underresponsive/Seeks Sensation, in which Typical Performance was found only in 39.2%, 42.2% and 46.1%, respectively. On the contrary, there was less impairment in Taste/Smell Sensitivity, Auditory Filtering and Visual/Auditory Sensitivity, in which Typi-

cal Performance was found in 81.4%, 64.7% and 76.5%, respectively.

Be that as it may, this study did not include individuals with ASD as a control group. It is speculated that the individuals with PWS in this study were not as severe as those with ASD in terms of the levels of sensory processing ability. This possibility is suggested by the above mentioned study conducted by Tomchek and Dunn⁵⁰⁾. Based on the total score of the SSP, they found 83.6% of the individuals with ASD were classified into the group of Definite Difference, 11.4% into Probable Difference and only 5.0% into Typical Performance.

In regards to sensory processing, this study failed to find significant differences between DEL and mUPD genetic subtypes, and those between male and female. The sole exception was gender difference in Underresponsive/Seek Sensation section, in which male individuals were more severely impaired than female individuals. Nonetheless, these findings obtained by rough analyses should be interpreted with caution, because participants in this study did not sufficiently control confounding factors, such as age, intelligence, BMI and complications. Likewise, a precise probe into each of the seven subscores should be carried out based on sufficient sample size. Despite statistically insignificant, the *p*-value ($p = 0.06$) of




Fig. 1 The effects of the SSP-J total score on the total score and subscores of ABC-J
 * $p < 0.05$, ** $p < 0.01$.

Auditory Filtering in inter-genotypical comparison was exceptionally low, as compared with those of the other subscores ($p = 0.37 - 0.98$). Due to the small sample size, type II error might conceal the possibility that mUPD showed a more severe impairment than DEL in terms of Auditory Filtering.

As far as the relationship of sensory processing with other behavioral symptoms was concerned, aberrant behaviors brought out a sharp contrast with food-related problems. No significant relationship was found between sensory processing and food-related behaviors. On the contrary, significant associations were observed between sensory processing and aberrant behaviors. Such a finding held true across diverse types of aberrant behaviors, including excitement, apathy, stereotype, hyperactivity and inappropriate speech. Regardless of differences in behavioral manifestations, it was observed that sensory processing abnormalities were highly linked with maladaptive behaviors. Similar findings supporting a predictive association between sensory processing dysfunction

and problem behaviors have been demonstrated in ASD^{51,52}. In this respect, the current study is the first report to confirm the significant relationship between sensory processing dysfunction and problem behaviors in PWS.

Questions remain unanswered whether sensory processing impairments are a component of core behavioral symptoms or the concomitant phenomenon. Even in ASD, a debate is still ongoing whether sensory processing deficits are an essential attribute or an accidental property⁵³. A systematic aggregation of evidence is needed to clarify whether sensory symptoms should be regarded as core behavioral features of PWS (i.e. temper tantrums, compulsive, ritualistic behaviors, skin picking behaviors and autistic-like behaviors).

This study found lack of significant relationship between food-related behaviors and sensory processing impairments in PWS. Such a pattern was found in maladaptive food-related behaviors, such as impairment of satiety, preoccupation with food and other

food-related negative behaviors. In ASD, on the other hand, relationships have been found between variables relating to food intake and those relating to sensory reactivity. For example, Tanner et al.⁵⁴⁾ demonstrated a positive relationship between total food eaten and SSP Taste/Smell Sensitivity scores, and a negative relationship between limited food variety and SSP Taste/Smell Sensitivity scores. As it has already been pointed out⁵⁵⁾, abnormal eating behaviors in PWS may be derived from a dysfunction in a satiety system, but not in the form of hunger. It is suspected that sensory processing impairments do not play a significant role in the mechanisms underlying hyperphagia related to lack of satiety in PWS.

This study has some methodological limitations. First, the effect of chronological age and intelligence was not fully considered. Unfortunately, the number of participants in this study was too small to analyze sensory profile differences in multiple age and/or intellectual groups. Second, this study failed to assess the influence of endocrinological factors including growth hormone therapy, obesity and diabetes. As there is some evidence to suggest the negative impact of type 2 diabetes on sensory processing⁵⁶⁾, the comparison between PWS individuals with and those without diabetes in regard to sensory responsiveness is of particular interest. Third, the impacts of other complications (scoliosis, cellulitis, etc.) were not examined. It has been well known that individuals with PWS have an incidence of scoliosis at rates between 40–90%^{57,58)}. At the same time, orthopedics literature has often pointed out that individuals with PWS have an increased tolerance to pain, which may potentially be helpful in the stage of rehabilitation after surgery⁵⁹⁾. Equally well known is the fact that individuals with PWS often engage in self-harming behaviors, such as compulsive skin picking and gouging. Consequently, erysipelas and cellulitis are common skin complications associated with PWS. Future research is needed to investigate the relationship between sensory processing differences and related problematic behaviors, considering the influence of orthopedic and dermatological complications.

Data Availability Statement

The datasets used to support the findings of this

study are available from the corresponding author on reasonable request.

Conflict of Interests

None

Acknowledgements. This research was supported by a Health Labor Science Research Grant in the fiscal year 2018 (Heisei 30).

REFERENCES

- 1) Dunn W, Brown C : Factor analysis on the Sensory Profile from a national sample of children without disabilities. *Am J Occup Ther* **51** : 490–495, 1997.
- 2) John AE, Mervis CB : Sensory modulation impairments in children with Prader–Willi syndrome. *Am J Med Genet C Semin Med Genet* **154C** : 266–276, 2010.
- 3) Gal E, Cermak SA, Ben–Sasson A : Sensory processing disorder in children with autism. *Growing Up with Autism : Working with School–Age Children and Adolescents*, New York, Guilford Press, pp95–123, 2007.
- 4) Mangeot SD, Miller LJ, McIntosh DN, et al : Sensory modulation dysfunction in children with attention-deficit–hyperactivity disorder. *Dev Med Child Neurol* **43** : 399–406, 2001.
- 5) Baranek GT, Chin YH, Hess LM, et al : Sensory processing correlates of occupational performance in children with fragile X syndrome : preliminary findings. *Am J Occup Ther* **56** : 538–546, 2002.
- 6) Riby DM, Janes E, Rodgers J : Brief report : exploring the relationship between sensory processing and repetitive behaviours in Williams Syndrome. *J Autism Dev Disord* **43** : 478–482, 2013.
- 7) Hilton CL, Harper JD, Kueker RH, et al : Sensory responsiveness as a predictor of social severity in children with high functioning autism spectrum disorders. *J Autism Dev Disord* **40** : 937–945, 2010.
- 8) Prader A, Labhart A, Willi H, et al : Ein Syndrom von Adipositas, Kleinwuchs, Kryptorchismus und Idiotie bei Kindern und Erwachsenen, die als Neugeborene ein myotonieartiges Bild geboten haben. VIII International Congress of Paediatrics, Copenhagen, 1956.
- 9) Crinò A, Schiaffini R, Ciampalini P, et al : Hypogonad-

- ism and pubertal development in Prader-Willi syndrome. *Eur J Pediatr* **162** : 327-333, 2003.
- 10) Buiting K, Saittoh S, Gross S, et al : Inherited microdeletions in the Angelman and Prader-Willi syndromes define an imprinting centre on human chromosome 15. *Nat Genet* **9** : 395-400, 1995.
 - 11) Cassidy SB, Driscoll DJ : Prader-Willi syndrome. *Eur J Hum Genet* **17** : 3-13, 2009.
 - 12) Nicholls RD, Saitoh S, Horsthemke B : Imprinting in Prader-Willi and Angelman syndromes. *Trends Genet* **14** : 194-200, 1998.
 - 13) Ledbetter DH, Riccardi VM, Airhart SD, et al : Deletions of chromosome 15 as a cause of the Prader-Willi syndrome. *N Engl J Med* **304** : 325-329, 1981.
 - 14) Holland AJ, Treasure J, Coskeran P, et al : Measurement of excessive appetite and metabolic changes in Prader-Willi syndrome. *Int J Obes Relat Metab Disord* **17** : 527-532, 1993.
 - 15) Tunncliffe P, Woodcock K, Bull L, et al : Temper outbursts in Prader-Willi syndrome : causes, behavioural and emotional sequence and responses by carers. *J Intellect Disabil Res* **58** : 134-150, 2014.
 - 16) Descheemaeker MJ, Vogels A, Govers V, et al : Prader-Willi syndrome : new insights in the behavioural and psychiatric spectrum. *J Intellect Disabil Res* **46** : 41-50, 2002.
 - 17) Dykens EM, Leckman JF, Cassidy SB : Obsessions and compulsions in Prader-Willi syndrome. *J Child Psychol Psychiatry* **37** : 995-1002, 1996.
 - 18) Greaves N, Prince E, Evans DW, et al : Repetitive and ritualistic behaviour in children with Prader-Willi syndrome and children with autism. *J Intellect Disabil Res* **50** : 92-100, 2006.
 - 19) Arron K, Oliver C, Moss J, et al : The prevalence and phenomenology of self-injurious and aggressive behaviour in genetic syndromes. *J Intellect Disabil Res* **55** : 109-120, 2011.
 - 20) Klabunde M, Saggat M, Hustyi KM, et al : Neural correlates of self-injurious behavior in Prader-Willi syndrome. *Hum Brain Mapp* **36** : 4135-4143, 2015.
 - 21) Descheemaeker MJ, Govers V, Vermeulen P, et al : Pervasive developmental disorders in Prader-Willi syndrome : the Lenven experience in 59 subjects and controls. *Am J Med Genet A* **140** : 1136-1142, 2006.
 - 22) Dykens EM, Lee E, Roof E : Prader-Willi syndrome and autism spectrum disorders : an evolving story. *J Neurodev Disord* **3** : 225-237, 2011.
 - 23) Dimitropoulos A, Schultz RT : Autistic-like symptomatology in Prader-Willi syndrome : a review of recent findings. *Curr Psychiatry Rep* **9** : 159-164, 2007.
 - 24) Milner KM, Craig EE, Thompson RJ, et al : Prader-Willi syndrome : intellectual abilities and behavioural features by genetic subtype. *J Child Psychol Psychiatry* **46** : 1089-1096, 2005.
 - 25) Veltman MW, Thompson RJ, Roberts SE, et al : Prader-Willi syndrome--a study comparing deletion and uniparental disomy cases with reference to autism spectrum disorders. *Eur Child Adolesc Psychiatry* **13** : 42-50, 2004.
 - 26) Bolton PF, Dennis NR, Browne CE, et al : The phenotypic manifestations of interstitial duplications of proximal 15q with special reference to the autistic spectrum disorders. *Am J Med Genet* **105** : 675-685, 2001.
 - 27) Veltman MW, Craig EE, Bolton PF : Autism spectrum disorders in Prader-Willi and Angelman syndromes : a systematic review. *Psychiatr Genet* **15** : 243-254, 2005.
 - 28) Hogart A, Wu D, LaSalle JM, et al : The comorbidity of autism with the genomic disorders of chromosome 15q11.2-q13. *Neurobiol Dis* **38** : 181-191, 2010.
 - 29) Kern JK, Trivedi MH, Garver CR, et al : The patterns of sensory processing abnormalities in autism. *Autism*, **10** : 480-494, 2006.
 - 30) Case-Smith J, Weaver LL, Fristad MA : A systematic review of sensory processing interventions for children with autism spectrum disorders. *Autism* **19** : 133-148, 2015.
 - 31) Marco EJ, Hinkley LB, Hill SS, et al : Sensory processing in autism : a review of neurophysiologic findings. *Pediatr Res* **69** : 48R-54R, 2011.
 - 32) Haesen B, Boets B, Wagemans J : A review of behavioural and electrophysiological studies on auditory processing and speech perception in autism spectrum disorders. *Research in Autism Spectrum Disorders* **5** : 701-714, 2011.
 - 33) Simmons DR, Robertson AE, McKay LS, et al : Vision in autism spectrum disorders. *Vision Res* **49** : 2705-2739, 2009.
 - 34) Tavassoli T, Baron-Cohen S : Taste identification in adults with autism spectrum conditions. *J Autism*

- Dev Disord **42** : 1419-1424, 2012.
- 35) Hilton CL, Graver K, LaVesser P : Relationship between social competence and sensory processing in children with high functioning autism spectrum disorders. *Research in Autism Spectrum Disorders* **1** : 164-173, 2007.
 - 36) American Psychiatric Association : Diagnostic and statistical manual of mental disorders, 5th edition. Arlington, VA, 2013.
 - 37) Ben-Sasson A, Carter AS, Briggs-Gowan MJ : Sensory over-responsivity in elementary school : prevalence and social-emotional correlates. *J Abnorm Child Psychol* **37** : 705-716.
 - 38) Wechsler D : Wechsler Intelligence Scale for Children-3rd ed. The Psychological Corporation, Sun Antonio, TX, 1991.
 - 39) Wechsler D : Wechsler Adult Intelligence Scale-3rd ed. The Psychological Corporation, San Antonio, TX, 1997.
 - 40) Japanese WISC- III Publication Committee : Nihonban WISC- III chinou kensahou (Japanese Wechsler Intelligence Scale for Children-3rd ed). Nihon Bunka Kagakusya, 1998.
 - 41) Japanese WAIS- III Publication Committee : Nihonban WAIS- III chinou kensahou (Japanese Wechsler Adult Intelligence Scale, 3rd ed). Nihon Bunka Kagakusya, 2006.
 - 42) Tsuji M, Hagiwara T, Iwanaga R, et al : The Japanese version of sensory profile. Nihon Bunka Kagakusya, 2015.
 - 43) Dunn W : Sensory profile's user manual. The Psychological Corporation, San Antonio, TX, 1999.
 - 44) McIntosh DN, Miller LJ, Shyu V, et al : Short sensory profile. The Psychological Corporation, New York, 1999.
 - 45) Russell H, Oliver C : The assessment of food-related problems in individuals with Prader-Willi syndrome. *Br J Clin Psychol* **42** : 379-392, 2003.
 - 46) Aman MG, Singh NN, Ono Y : Clinical evaluation of Aberrant Behavior Checklist Japanese Version (ABC-J). Jiho, Tokyo, 2006.
 - 47) Shedlack KJ, Hennen J, Magee C, et al : A comparison of the Aberrant Behavior Checklist and the GAF among adults with mental retardation and mental illness. *Psychiatr Serv* **56** : 484-486, 2005.
 - 48) Brinkley J, Nations L, Abramson RK, et al : Factor analysis of the aberrant behavior checklist in individuals with autism spectrum disorders. *J Autism Dev Disord* **37** : 1949-1959, 2007.
 - 49) Schroeder SR, Rojahn J, Reese RM : Brief report : reliability and validity of instruments for assessing psychotropic medication effects on self-injurious behavior in mental retardation. *J Autism Dev Disord* **27** : 89-102, 1997.
 - 50) Tomchek SD, Dunn W : Sensory processing in children with and without autism : a comparative study using the short sensory profile. *Am J Occup Ther* **61** : 190-200, 2007.
 - 51) O'Donnell S, Deitz J, Kartin D, et al : Sensory processing, problem behavior, adaptive behavior, and cognition in preschool children with autism spectrum disorders. *Am J Occup Ther* **66** : 586-594, 2012.
 - 52) Lane AE, Young RL, Baker AE, et al : Sensory processing subtypes in autism : association with adaptive behavior. *J Autism Dev Disord* **40** : 112-122, 2010.
 - 53) Ben-Sasson A, Hen L, Fluss R, et al : A meta-analysis of sensory modulation symptoms in individuals with autism spectrum disorders. *J Autism Dev Disord* **39** : 1-11, 2009.
 - 54) Tanner K, Case-Smith J, Nahikian-Nelms M, et al : Behavioral and physiological factors associated with selective eating in children with autism spectrum disorder. *Am J Occup Ther* **69** : 6906180030 p1-p8, 2015.
 - 55) Hinton EC, Holland AJ, Gellatly MS, et al : Neural representations of hunger and satiety in Prader-Willi syndrome. *Int J Obes* **30** : 313-321, 2006.
 - 56) Cosway R, Strachan MW, Dougall A, et al : Cognitive function and information processing in type 2 diabetics. *Diabet Med* **18** : 803-810, 2001.
 - 57) Holm VA, Lurnen EL : Prader-Willi syndrome and scoliosis. *Dev Med Child Neurol* **23** : 192-201, 1981.
 - 58) Nagai T, Obata K, Ogata T, et al : Growth hormone therapy and scoliosis in patients with Prader-Willi syndrome. *Am J Med Genet A* **140** : 1623-1627, 2006.
 - 59) Van-Bosse HJP : Guidelines on scoliosis monitoring and treatment for children with Prader-Willi syndrome. Prader-Willi Syndrome Association, Florida, 2010.