

**Table 4 Comparison of in-hospital events within 30 days post-admission between low- and high-risk groups based on ER French Calculator**

	Low-risk group (n=29)	High-risk group (n=23)	P
Embolic complications within 30 days			
Total; n (%)	5 (12)	10 (45)	0.0381
Cerebral; n (%)	4 (9)	8 (36)	0.0744
Renal; n (%)	0 (0)	1 (4.5)	0.2569
Cerebro-renal; n (%)	0 (0)	1 (4.5)	0.2569
Peripheral; n (%)	1 (3)	0 (0)	0.3685
Surgical treatment within 30 days; n (%)			
Prevention of embolic events; n (%)	4 (14)	9 (39)	0.0015
Primary prevention; n (%)	1 (3)	2 (9)	0.2439
Secondary prevention; n (%)	3 (10)	7 (30)	0.0679
Uncontrolled infection; n (%)	1 (3)	6 (26)	0.0175
Refractory heart failure; n (%)	0 (0)	2 (9)	0.1054
Death within 30 days; n (%)	0 (0)	1 (4.5)	0.2569
Composite of events within 30 days; n (%)	7 (24)	21 (91)	<0.0001

Low-risk group: probability on the 28<sup>th</sup> day  $\leq$ 8%, High-risk group, probability on the 28<sup>th</sup> day  $>$ 8%

