

High-density Lipoprotein Cholesterol Efflux Capacity as a Novel Prognostic Surrogate for
Coronary Artery Disease

Itaru Hisauchi¹, Tetsuya Ishikawa¹, Makoto Ayaori², Harumi Uto-Kondo^{2,3}, Yuri Koshikawa¹,
Tomoaki Ukaji¹, Hidehiko Nakamura¹, Yukiko Mizutani¹, Isao Taguchi¹, Takatomo
Nakajima⁴, Makoto Mutoh⁴, Katsunori Ikewaki²

¹Department of Cardiology, Dokkyo Medical University, Saitama Medical Center, Koshigaya,
Saitama, Japan

²Division of Anti-aging and Vascular Medicine, Department of Internal Medicine, National
Defense Medical College, Tokorozawa, Saitama, Japan

³Department of Bioscience in Daily Life, Nihon University, Fujisawa, Kanagawa, Japan

⁴Division of Cardiology, Saitama Cardiovascular Respiratory Center, Kumagaya, Saitama,
Japan

Corresponding author: Tetsuya Ishikawa, MD, PhD

Tel.: +81-485-36-9900 ext. 2250

FAX: +81-485-36-9900

E-mail: tetsuya50ishikawa@gmail.com

Running head: Cholesterol Efflux Capacity and MACE

COI: All of the authors have no conflict of interest, financial or otherwise.

Abstract

Aim: We examined the impact of baseline high-density lipoprotein cholesterol efflux capacity (CEC) on major cardiac adverse events (MACE) in patients with coronary artery disease (CAD) during a long-term secondary prevention.

Method: CEC was measured using a cell-based efflux system in (3)[H]-cholesterol-labeled J774 macrophages in apolipoprotein B-depleted plasma between January 2011 and January 2013. Patients with CAD were divided into 2 groups as a boundary CEC value of 1: $0.19 \leq \text{CEC} < 1$ (impaired CEC group, mean CEC of 0.76 ± 0.16 , $n = 136$), and $1 \leq \text{CEC} \leq 2.08$ (enhanced CEC group, 1.20 ± 0.19 , $n = 44$). MACE, comprised the incidence of cardiac death, non-fatal myocardial infarction, and any revascularizations (RV) without restenosis approximately 1 year after vascularization, was retrospectively investigated at September 2019. Impact of enhanced CEC on MACE among 22 variables was examined by applying a Cox proportional hazard model.

Result: The frequency of MACE in impaired CEC group (16.9%, mean observational interval of 2111 ± 888 days) was significantly higher than that in enhanced CEC group (2.3%, 2252 ± 685 , $p = 0.013$), largely driven by the significantly higher RV incidence (14.0 % versus 2.3 %,

p = 0.032). Enhancement of CEC was the significant predictor of MACE (hazard ratio: 0.11; 95% CI: 0.013-0.879; p = 0.038).

Conclusion: A baseline CEC level of more than 1 in patients with CAD brought favorable long-term clinical outcomes, suggesting that CEC is a useful prognostic and therapeutic surrogate for secondary prevention of CAD.

247 words

Keywords: Coronary artery disease, high-density lipoprotein cholesterol efflux capacity, major adverse cardiac event, coronary revascularization

Introduction

In addition to low-density lipoprotein cholesterol (LDL-C), it has been essential to clarify both the prognostic surrogate and the therapeutic target for secondary prevention in patients with coronary artery disease (CAD); this is because relative risks of secondary cardiac events remained despite the lowering therapy of LDL-C [1]. Among the pleiotropic atheroprotective effects of high-density lipoprotein cholesterol (HDL-C), cholesterol efflux from macrophage (cholesterol efflux capacity, CEC) was consistently demonstrated as a negative predictor of CAD, independent of HDL-C and LDL-C levels [2,3]. In addition, the CEC in asymptomatic healthy subjects predicts the incidence of atherosclerotic cardiovascular events (ASCVD) corresponding to the baseline CEC level besides the baseline LDL-C level [4]. However, the prognostic impact of the baseline CEC on patients with CAD during long-term secondary prevention therapy is not fully understood.

Therefore, we sought to clarify the impact of CEC on cardiac events during secondary prevention in patients with CAD in the present drug-eluting stent era. In order to examine the relationship between the baseline CEC levels and the incidences of major adverse cardiac events (MACE), we retrospectively investigated the frequency of MACE by dividing 180 patients with CAD enrolled in our previous study [3] into 2 groups with the border baseline

CEC value of 1.0, the normal function of CEC. The baselines and the frequencies of clinical outcomes were compared between the two groups, i.e. impaired CEC group (CEC less than 1) versus enhanced CEC group (CEC more than 1). In addition, the impact of enhancement of CEC on MACE among 22 variables was examined by using a Cox proportional hazard model.

Methods

Population

The current study was an observational study that investigated the relationship between the baseline CEC levels and the clinical outcomes in 180 patients with CAD corresponding to our previous report [3]; there were two cases of in-hospital mortality among the CAD patients in our previous report [3]. Fasting blood sampling was performed at the time of elective CAG, elective percutaneous coronary intervention (PCI), or multi-slice coronary computed tomography (MSCT) between January 2011 and January 2013 at Saitama Cardiovascular Respiratory Center (SCRC). All the patients had continuous secondary prevention according to the guideline of the Japanese Circulation Society [5]. All patients were informed of the study objectives, and consent was obtained from all participants. The rationale of the present study was approved by the local ethics committee of SCRC on 14, February 2019 (accepted number: 2018040). Retrospective investigation of medical records from the SCRC, and telephone and letter interviews of the clinics and patients were conducted from March 2019 to September 2019.

Based on the baseline levels of CEC, patients were divided into two groups as the border value of 1, the normal range of CEC: $0.19 \leq \text{CEC} < 1.0$ (impaired CEC group, $n = 136$), and $1.0 \leq \text{CEC} \leq 2.08$ (enhanced CEC group, $n = 44$).

Measurement of cholesterol efflux capacity of macrophages

The CEC was determined according to the methods previously reported [2,3]. In brief, J774 macrophages were purchased from RIKEN (Tsukuba, Japan), cultured in RPMI 1640 medium containing 10% fetal bovine serum, and kept under constant conditions of 5% carbon dioxide and a temperature of 37°C. J774 cells were plated in 24-well plates, grown to 80% confluence, and radiolabeled with 2 $\mu\text{Ci/mL}$ of ^3H -cholesterol. Apolipoprotein B-depleted serum was prepared by incubation with 13% polyethylene glycol 6000 solution (Wako Pure Chemicals). Subsequently, an efflux medium containing 2.8% apolipoprotein B-depleted serum was added and incubated for 4 h. All procedures were performed in the presence of the acyl-coenzyme A: cholesterol acyltransferase inhibitor Sandoz 58-035 (2 $\mu\text{g/mL}$; Sigma, St. Louis, MO, USA) and 8-bromoadenosine 3', 5'-cyclic monophosphate (0.3 mmol/L; Sigma). A liquid scintillation counter was used to quantify the efflux of radioactive cholesterol from the cells. The quantity of radioactive cholesterol incorporated into cellular lipids was calculated through hexane:isopropanol (*v:v*, 1:1) extraction in control wells not exposed to the serum. The percent

efflux was calculated using the following formula: $(\text{cpm of } ^3\text{H-cholesterol in media containing 2.8\% apolipoprotein B-depleted serum} - \text{cpm of } ^3\text{H-cholesterol in serum-free mediums}) / (\text{cpm of } ^3\text{H-cholesterol in cells extracted before the efflux step}) \times 100$. All assays were performed in duplicate. The CECs of patients' sera were expressed as the values relative to those of the pooled sera.

Baseline measurements

Baseline variables were collected from our previous report [3]. The patient variables included age, male sex, and the coronary risk factors such as hypertension, dyslipidemia, diabetes mellitus (diabetes), and family history of CAD. Patients with no history of smoking (nonsmoker) were included as the risk factor. Patients with hypertension were considered to be at risk if their blood pressure was $\geq 140 / 90$ mm Hg or if they had a history of anti-hypertensive drug use. Patients with diabetes were considered to be at risk if their fasting glucose level was ≥ 126 mg/dL and their hemoglobin A1C (HbA1c) level was $\geq 6.5\%$, or if they had a history of hypoglycemic drug or insulin use. Patients with dyslipidemia were considered to be at risk if their LDL-C level was ≥ 140 mg/dL and their HDL-C level was ≤ 40 mg/dL, or if they were taking a lipid-lowering drug, including HMG-CoA reductase inhibitors (statins). Dyslipidemia-related variables, including serum total cholesterol, triglycerides,

HDL-C, LDL-C, and CEC were evaluated. Apolipoprotein (apo) A-I (apo-A1) and apo-B concentrations were measured by turbidimetric immunoassay. The administration of statins (statin administration) was not prospectively randomized, and patients who were prescribed statins at the time of blood sampling were evaluated as taking statins based on the physician's discretion. The cardiovascular baselines, serum hematocrit (Ht), creatinine (Cr), and brain natriuretic peptide (BNP) levels were evaluated. The percentage of left ventricular ejection fractions less than 40% (left ventricular dysfunction), prevalence of coronary artery bypass grafting (CABG), diseased left main coronary artery, and prevalence of peripheral artery disease (PAD) were also estimated, as were the mean numbers of diseased coronary vessels (number of diseased coronary vessel).

The estimated endpoint

The incidence of major adverse cardiac events (MACE) comprised the incidences of cardiac death, including sudden death, non-fatal myocardial infarction, and any target lesion revascularization (TLR) without restenosis, approximately 1 year after vascularization. The incidence of TLR was divided into late TLR, defined as TLR at the previous target lesion after 1-year secondary angiogram [6], and non-culprit TLR, defined as the TLR for de novo lesions without culprit revascularized lesions [7]. All of the late TLR was conducted against the

failure of the first-generation drug-eluting stents (i.e. sirolimus-eluting stent placement, and paclitaxel-eluting stents) placed prior to February 2010.

In addition, the incidence of all-cause mortality was investigated as the other estimated outcome. The clinical observational interval was the interval from the date of blood sampling of CEC to the date of the final confirmation of the clinical course.

Statistical analyses

The 23 baseline characteristic variables were expressed as percentages or mean values \pm standard deviation. Baseline variables (Table 1) and clinical outcomes (Table 2) in impaired CEC group were individually compared with those in enhanced CEC group using unpaired *t*-tests for continuous values and the χ^2 test for categorical values. The cumulative MACE-free ratio of impaired CEC group was compared with that of enhanced CEC group by log-rank test (Figure 1). The predictors of MACE were examined by applying a Cox proportional hazard model among 22 variables. The CEC was enrolled as the categorical group, i.e. enhanced CEC group. Dyslipidemia was excluded from this statistics because of the duplication of the individual dyslipidemia-related variables. A *p* value < 0.05 was considered statistically significant. The Stata version 14 software for Windows (StataCorp, College Station, TX, USA) was used for all statistical analyses.

Results

The baseline characteristics of the study subjects are shown in Table 1. None of the patient-related baselines and coronary risk factors in impaired CEC group was significantly different from those in enhanced CEC group. Among the dyslipidemia-related variables, the mean serum HDL-C, apo-A1, and CEC in impaired CEC group were significantly different from those in enhanced CEC group (HDL-C level: 48.2 ± 12.7 vs. 55.5 ± 15.9 mg/dL; $p = 0.006$; apo-A1 level: 120 ± 25.1 vs. 129 ± 25.1 mg/dL; $p = 0.039$; CEC: 0.76 ± 0.16 vs. 1.20 ± 0.19 ; $p < 0.001$). None of other laboratory variables in impaired CEC group was significantly different from those in enhanced CEC group. Among the cardiovascular-related baselines, the number of diseased coronary vessel in impaired CEC group was significantly larger than that in enhanced CEC group (1.98 ± 0.80 vs. 1.68 ± 0.71 ; $p = 0.018$).

The clinical outcomes are shown in Table 2. The frequencies of MACE, TLR, and non-culprit TLR in impaired CEC group were significantly higher than those in enhanced CEC group (16.9% vs. 2.3%; $p = 0.013$, 14.0% vs. 2.3%; $p = 0.032$, 9.6% vs. 0; $p = 0.033$, respectively).

The cumulative MACE-free ratios are shown in Figure 1. The cumulative MACE-free ratio of impaired CEC group was significantly lower than that in enhanced CEC group ($p = 0.022$, log-rank test).

The predictors of MACE in the entire cohort are shown in Table 3. The PAD (hazard ratio [HR]: 32.3; 95% confidence interval [CI]: 4.60-226; $p < 0.001$), and enhanced CEC (HR: 0.11; 95% CI: 0.013-0.879; $p = 0.038$) were the significant predictors of MACE (Table 3).

Discussion

The primary purpose of the present study was to examine the impact of the baseline CEC level of patients with CAD, particularly, the impact of the border CEC value of 1, on major adverse cardiac events (MACE) during long-term secondary prevention. As the backgrounds, in addition to the LDL-C level, it is essential to clarify the prognostic surrogate and/or the therapeutic target of secondary prevention in patients with CAD [1]. The diagnostic impact of CEC on CAD has been consistent [2,3], and the prognostic impact of CEC on ASCVD has been previously documented among asymptomatic healthy subjects [4]. To the best of our knowledge, the present study is the first to demonstrate the prognostic impact of baseline CEC level, but not HDL-C or apo-A1 levels, on MACE in patients with CAD with the longest clinical observational interval. The frequency of MACE was quite different at a border CEC level of 1 with the large reduction of the frequency of non-culprit TLR (the development of de novo lesion) in enhanced CEC group, supporting the atheroprotective effects of CEC against the atherosclerotic coronary stenosis [2,3,4]. Therefore, the present study showed 1) the prognostic impact of baseline CEC on long-term clinical outcomes during secondary prevention of CAD, 2) a useful therapeutic goal of CEC level of 1 for secondary prevention of

CAD, and 3) CEC as the single significant predictor of MACE in patients with CAD among conventional lipid-related variables.

The long-term clinical outcome in the cohort with CEC more than 1 (enhanced CEC group) was favorable in a few percentages of MACE (only one case of late TLR) during a mean clinical observation interval of more than 6 years (Table 2, Figure 1). The present low frequency of MACE in enhanced CEC group was similar to that of ASCVD at approximately 6 years follow-up in the cohort with CEC more than 1 in healthy young subjects [4]. However, in enhanced CEC group, there were several adverse baselines for MACE, such as a higher tendency of mean age than impaired CEC group [8], percentage of diabetes [9] as high as 50%, mean baseline BNP level more than 150 (pg/dL), and several percentages of the prevalence of PAD [10] (Table 1). All of the significantly higher CEC, HDL-C, and apo-A1 levels in enhanced CEC group than those of impaired CEC group were the atheroprotective variables (Table 1). However, CEC remained to be the significant predictor of MACE by a Cox proportional hazard model (Table 3). Thus, CEC was the single significant predictor of MACE in patients with CAD among not only lipid-related variables, but also other conventional coronary risk factors. Therefore, the present favorable clinical outcome of enhanced CEC group is dependent upon the significantly higher CEC. As described above, the TLR was inversely related to CEC (Table 2). Several predictors of late TLR after sirolimus-eluting stent

placement in patient, lesion, and procedure characteristics were reported (j-Cypher Registry) [6]. In addition, several predictors related to atherosclerotic coronary plaque for non-culprit TLR were reported (PROSPECT study) [11]. Thus, the atheroprotective effects of CEC played a significant role for, particularly, reducing the non-culprit TLR, i.e. development of de novo stenotic lesions (Table 2). Accordingly, the baseline CEC predicts the incidence of MACE, in particular, of coronary revascularization, in patients with CAD, including the boarder baseline CEC level of 1. Several interventions recommended as the optimal secondary preventions of CAD have been described and function to improve CEC level up to 1; these include pharmacological interventions by rosuvastatin [12], ezetimibe [13], eicosapentaenoic acid [14], and intensive cardiac rehabilitation, including smoking cessation [15] and improvement of diabetes state [9].

On the other hand, the frequency of MACE in impaired CEC group was as high as 16.9% in a mean observational interval of 5.78 years (approximately 7.3 fold higher compared to enhanced CEC group, Table 2, Figure 1). The present frequency of MACE in impaired CEC group was similar to the Japanese large-scale multicenter RESET trial (approximately 20% to 23% in a 7-year follow-up) [16]. This frequency of MACE was several times higher than that of ASCVD, which ranged from approximately 2% to 4% at a 6-year clinical follow-up in the relatively lower values of CEC in healthy young subjects [4]. Therefore, as described above,

the CEC level of 1, the boarder of the normal range of CEC, exerted an important prognostic impact on MACE in patients with CAD. In the present patients with CAD, PAD predicts the incidence of MACE beyond baseline CEC level (Table 3), although the percentages of the patient with PAD were approximately 5% (Table 1). Patients with CAD concurrent with PAD have a high risk for MACE, and are recommended intensive lipid lowering treatment [10]. Accordingly, in the CAD cohorts with CEC level less than 1, the impact of CEC on MACE with further intensive lipid-lowering therapy should be examined.

The present study had several limitations; these included a small population, retrospective observational study, factors related to the coronary stenotic lesion and the PCI procedure, achievement of CEC after secondary prevention, and the impact of other factors and underlying confounders potentially related to MACE during a long-term interval.

Conclusion

The present study demonstrated that 1) the baseline CEC level more than 1 in patients with CAD brought favorable long-term clinical outcomes, 2) CEC is a useful prognostic surrogate for the secondary prevention of CAD, and 3) CEC is the single significant predictor of MACE among not only lipid-related variables, but also conventional coronary risk factors.

Acknowledgments and Notice of grant support

This study was supported by a Research Fund from Saitama Cardiovascular and Respiratory Center under Grants 10DR, 11DR, 12DR, and 13DR, and Grants-in-aid for Scientific Research for T.I. (18H02806).

References

1. Robinson JG, Farnier M, Krempf M, et al. ; ODYSSEY LONG TERM Investigators. Odyssey long term investigators. Alirocumab and cardiovascular outcomes after acute coronary syndrome. *N Engl J Med.* (2015); 372: 1489-99.
2. Khera AV, Cuchel M, de la Llera-Moya M, et al. Cholesterol efflux capacity, high-density lipoprotein function, and atherosclerosis. *N Engl J Med.* 2011, Jan 13;364(2):127-35.
3. Ishikawa T, Ayaori M, Uto-Kondo H, et al. High-density lipoprotein cholesterol efflux capacity as a relevant predictor of atherosclerotic coronary disease. *Atherosclerosis.* (2015); 242: 318-22.
4. Rohatgi A, Khera A, Berry JD, et al. HDL cholesterol efflux capacity and incident cardiovascular events. *N Engl J Med.* 2014 Dec 18;371(25):2383-93.
5. Ishikawa K, et al. Guidelines for the Diagnosis and Treatment of Cardiovascular Diseases (2004–2005 Joint Working Groups Report). Guidelines for Secondary Prevention of Myocardial Infarction (JCS 2006). http://www.j-circ.or.jp/guideline/pdf/JCS2006_ishikawa_h.pdf (in Japanese).
6. Nakagawa Y, Kimura T, Morimoto T, et al; j-Cypher Registry Investigators. Incidence and risk factors of late target lesion revascularization after sirolimus-eluting stent implantation (3-year follow-up of the j-Cypher registry). *Am J Cardiol.* (2010) 106: 329-336.

7. Stone GW, Maehara A, Lansky AJ, et al; PROSPECT Investigators. A prospective natural-history study of coronary atherosclerosis. *N Engl J Med.* (2011) 364: 226-235.
8. Berrougui H, Khalil A. Age-associated decrease of high-density lipoprotein-mediated reverse cholesterol transport activity. *Rejuvenation Res.* (2009) 12:117-126. doi: 10.1089/rej.2008.0840.
9. Zhou H, et al. Zhou H, Tan KC, et al. Cellular cholesterol efflux to serum is impaired in diabetic nephropathy. *Diabetes Metab Res Rev.* (2008) 24:617-23.
10. Jukema JW, Szarek M, Zijlstra LE et al ; ODYSSEY OUTCOMES Committees and Investigators. Alirocumab in patients with polyvascular disease and recent acute coronary syndrome: ODYSSEY OUTCOMES Trial. *J Am Coll Cardiol.* (2019) 74: 1167-1176.
11. BoZhengMD, Gary S.MintzMD, John A.McPhersonMD et al. <https://doi.org/10.1016/j.jcmg.2015.06.014> Predictors of plaque rupture within nonculprit fibroatheromas in patients with acute coronary syndromes. The PROSPECT study. *J Am Coll Cardiol Img.* (2015) 8: 1180-1187.
12. Kim KM, Jung KY, Yun HM, et al. Effect of Rosuvastatin on cholesterol efflux capacity and endothelial function in type 2 diabetes mellitus and dyslipidemia. *Circ J.* (2018) 82: 1387-1395.
13. Uto-Kondo H, Ayaori M, Sotherden GM, et al. Ezetimibe enhances macrophage reverse cholesterol transport in hamsters: contribution of hepato-biliary pathway. *Biochim Biophys Acta.* (2014); 1841: 1247-55.

14. Tanaka N, Ishida T, Nagao M, et al. Administration of high dose eicosapentaenoic acid enhances anti-inflammatory properties of high-density lipoprotein in Japanese patients with dyslipidemia. *Atherosclerosis*. (2014); 237: 577-83.
15. Koba S, Ayaori M, Uto-Kondo H, et al. Beneficial effects of exercise-based cardiac rehabilitation on high-density lipoprotein-mediated cholesterol efflux capacity in patients with acute coronary syndrome. *J Atheroscler Thromb*. (2016); 23: 865-77.
16. Shiomi H, Kozuma K, Morimoto T, et al and on behalf of the RESET Investigators 7-Year Outcomes of a randomized trial comparing the first-generation sirolimus-eluting stent versus the new-generation everolimus-eluting stent: The RESET Trial. *JACC Cardiovasc Interv*. (2019) 12: 637-647.

Figure legends

Figure 1. Cumulative MACE-free ratios in impaired CEC group and enhanced CEC group

The cumulative MACE-free ratios in the 2 groups were determined by Kaplan-Meier curves.

The cumulative MACE-free ratio in impaired CEC group (broken line) was significantly lower than that in enhanced CEC group (solid line) ($p = 0.022$, log-rank test). The vertical axis shows the cumulative MACE-free ratio (%), while the horizontal axis shows the interval for clinical observation after CEC measurement (days).