

Original article**Evaluation of common carotid artery and carotid bulb plaque: A population-based study from Japan**

Short title: Factors influencing max-IMT in a Japanese population

Haruki Igarashi^{1,2}, Akio Iwasaki^{1,2}, Ryuta Okabe^{1,3}, Ayano Suzuki^{1,2}, Keisuke Suzuki¹,
Hidehiro Takekawa^{1,2,4}

1 Department of Neurology, Dokkyo Medical University, Tochigi, Japan

2 Stroke Center, Dokkyo Medical University, Tochigi, Japan

3 Department of Internal Medicine, Akiru Municipal Medical Center, Tokyo, Japan

4 Center of Medical Ultrasonics, Dokkyo Medical University, Tochigi, Japan

Corresponding author

Akio Iwasaki, M.D., Ph.D.

Department of Neurology, Dokkyo Medical University, 880 Kitakobayashi, Mibu, Tochigi

321-0293

Tel: +81-282-86-1111 (extension 2721), Fax: +81-282-86-5884.

e-mail: a.iwasaki.n@gmail.com

Abstract, **202** words; text, **3135** words; references, **34**; Tables, **3**; Figures, **none**;

Supplementary files, **none**.

Acknowledgments

We express our heartfelt thanks to Yuka Tsukahara, MD, PhD, and Madoka Okamura, MD, PhD, of the Department of Neurology, Dokkyo Medical University, for their assistance in conducting this study.

Conflict of interest

There are no financial or other relations that could lead to conflicts of interest.